ONE STEP DISINFECTANT

CLEANER - FUNGICIDE - MILDEWSTAT - VIRUCIDE* - DEODORIZER FOR HOMES, HOSPITALS, INSTITUTIONAL AND INDUSTRIAL USE ACTIVE INGREDIENTS.

VIRO-STAT RTU

bhard

ACTIVE INGREDIENTS.	
Octyl decyl dimethyl ammonium chloride	0.025%
Dioctyl dimethyl ammonium chloride	0.010%
Didecyl dimethyl ammonium chloride	0.015%
Alkyl (C14, 50%; C12, 40%; C16,10%) dimethyl benzyl ammonium chloride	0.034%
OTHER INGREDIENTS	
TOTAL	

KEEP OUT OF REACH OF CHILDREN

CAUTION **PRECAUTIONARY STATEMENTS** HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID: IF IN EYES: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. Have the container or label with you when calling a poison control center or doctor, or going for treatment.

> EPA Reg. No. 6836-152-11547 • EPA Est. No. 10088-WI-1 Net Contents ONE QUART (32 FL. 0Z.)

> > Designer:

VIRO-STAT RTU is a one-step Hospital use disinfectant, Bactericidal according to the current AOAC Use-Dilution Test Method, Eungicidal according to the AOAC Eungicidal Test and Vigucidal* according to the vigucidal gualification modified. in the presence of 5% organic serum against: (Bacteria:) Pseudomonas aeruginosa (Pseudomonas), Staphylococcus aureus (Staph), Salmonella enterica (Salmonella), Acinetobacter baumannii (Acinetobacter), Brevibacterium ammoniagenes (Brevibacterium), Campylobacter ieiuni (Campylobacter), Enterobacter aerogenes, Enterococcus faecalis (Enterococcus), Enterococcus faecalis - Vancomycin resistant (VRE), Escherichia coli (E. coli), ESBL Escherichia coli -(Extended spectrum beta-lactamase producing E, coli), Klebsiella pneumoniae (Klebsiella), Legionella pneumophilia, Pseudomonas cepacia (Pseudomonas), Salmonella schottmuelleri (Salmonella), Salmonella typhi (Salmonella), Serratia marcescens, Shigella dysenteriae (Shigella), Staphylococcus aureus Methicillin-Resistant (MRSA), Staphylococcus aureus Multi-Drug Resistant, Staphylococcus aureus - Vancomycin Intermediate Resistant - (VISA), Streptococcus pyogenes [Strep] [a cause of scarlet fever]. Vibrio cholerae. (Viruses:) *Hepatitis B Virus (HBV). * Hepatitis C Virus (HCV). *Herpes Simplex Virus Type 2, *HIV-1 (AIDS Virus), Human Coronavirus, *Influenza Virus Type A/Brazil (Influenza), *Norwalk Virus [Norovirus], *Respiratory Syncytial Virus (RSV), *SARS Associated Coronavirus (SARS) [cause of Severe Acute Respiratory Syndrome], *Vaccinia (Pox Virus). (Fungi:) Trichophyton mentagrophytes (the Athlete's Foot Fungus) (A Cause of Ringworm) (A Cause of Ringworm of the foot), Aspergillus niger,

Use VIRO-STAT RTU on washable hard, nonporous surfaces of: floors, walls, countertops, stovetops, sinks, exterior appliances, exterior refrigerators, exterior microwave ovens, cabinets, highchairs, garbage cans, refrigerated storage and display equipment, tables, non-wooden picnic tables, outdoor patio furniture except cushions and wood frames, chairs, desks, telephones, doorknobs, shower stalls, tubs, glazed tiles, kennel runs, cages, coils and drain pans of air conditioning and refrigeration equipment and heat pumps, conductive flooring and other hard nonporous surfaces made of: metal, stainless steel, glazed porcelain, glazed ceramic, sealed stone, plastic (such as polystyrene, polypropylene), glazed enameled surfaces, finished woodwork, vinyl and plastic upholstery.

DIRECTIONS FOR LISE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Before use in federally inspected meat and poultry food processing plants and dairies, food products and packaging materials must be removed from the room or carefully protected.

DISINFECTION/VIRUCIDAL*/FUNGICIDAL/MOLD & MILDEW CONTROL DIRECTIONS: Apply VIRO-STAT RTU undiluted to hard, nonporous surfaces, thoroughly wetting the surfaces with a cloth, mop, sponge, spraver or by immersion. Treated surfaces must remain wet for 10 minutes. Wipe dry with a cloth, sponge or mop or allow to air dry. For heavily soiled areas, a preliminary cleaning is recommended. For sprayer applications, use a coarse spray device. Spray 6 - 8 inches from surface; rub with brush, sponge or cloth. Avoid breathing spray. Rinse all surfaces that come in contact with food such as countertops, appliances, tables and stovetops with potable water before reuse. Do not use on utensils, glassware and dishes. For fungicidal activity: VIRO-STAT RTU is an effective one-step fungicide against Trichophyton mentagrophytes when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities. For mold and mildew: Repeat treatment every 7 days or more often if new growth appears. This product is not for use on medical device surfaces.

TO CLEAN AND DEODORIZE: Apply undiluted to surfaces with a cloth, mop, sponge or sprayer. Wipe or allow to air dry. KILLS HIV-1 and HBV and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/ BODY FLUIDS in health care settings or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS) or Hepatitis B Virus (HBV) or Hepatitis C Virus (HCV).

SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1 OR HBV OR HCV ON SURFACES/OBJECTS SOILED WITH BLOOD OR BODY FLUIDS. PERSONAL PROTECTION: Clean-up should always be done wearing protective gloves, gowns, masks and eve protection. CLEANING PROCEDURE: Blood and other body fluids containing HIV or HBV or HCV must be thoroughly cleaned from surfaces and objects before application of VIRO-STAT RTU.

CONTACT TIME: Leave surface wet for 1 minute (60 seconds) for HIV-1 and 10 minutes for HBV and HCV. Use a 10 minute contact time for disinfection against all other viruses, bacteria and fungi claimed. DISPOSAL OF INFECTIOUS MATERIAL: Blood, body fluids, cleaning materials and clothing should be autoclaved and disposed of according to local regulations for infectious waste disposal

(Animal Viruses:) Avian Infectious Bronchitis Virus, Canine Distemper Virus, Feline Calicivirus, Newcastle's Disease Virus, Pseudorabies Virus.

VETERINARY PRACTICE/ANIMAL CARE/ANIMAL LABORATORY/ZOOS/PET SHOP/KENNELS DISINFECTION/VIRUCIDAL* DIRECTIONS: For cleaning and disinfecting hard nonporous surfaces: equipment used for feeding or watering animals, utensils, instruments, cages, kennels, stables, catteries, pens, stalls and etc, Remove all animals and feeds from premises. animal transportation vehicles, crates, pens, stalls and etc. Remove all litter, droppings and manure from floors, walls and surfaces of facilities occupied or traversed by animals. Thoroughly clean all surfaces with soap or detergent and rinse with water. Empty all feeding and watering appliances. Saturate surfaces with VIRO-STAT RTU for a period of 10 minutes. Wipe or allow to air dry. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals as well as forks, shovels, and scrapers used for removing litter and manure. Thoroughly scrub all treated surfaces, then rinse all surfaces that come in contact with food, including equipment used for feeding or watering, with potable water before reuse. Ventilate buildings, animal enclosures, vehicles and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried

WATER DAMAGE RESTORATION AGAINST ODOR-CAUSING BACTERIA AND FUNGI FOR HOME, INSTITUTIONAL, INDUSTRIAL AND HOSPITAL USE: This product is particularly suitable for use in water damage restoration situations against odor causing bacteria on the following porous and semi-porous materials: carpets, carpet cushions, subfloors, drywall, trim and frame lumber, tackless strip and paneling. Saturate the affected materials with enough product to remain wet for at least 10 minutes. Sewer Backup and River Flooding: During mitigation procedures, remove gross filth or heavy soil along with non-salvageable materials. Saturate all affected areas with a coarse spray device before and after cleaning and extraction. Carpets, Carpet Cushions and Other Porous Materials such as Subfloors, Drywall, Trim and Frame Lumber, Tackless Strip and Paneling: For water damage from a clean source, extract excess water. Test hidden area for colorfastness. Remove gross filth or heavy soil. Saturate all affected areas with a coarse spray device. Roll, brush or agitate into the materials and allow materials to remain wet for 10 minutes. Follow with a thorough extraction. Dry rapidly and thoroughly. Special instructions For Cleaning Carpet Against Odor Causing Bacteria: Use this product in industrial and institutional areas such as homes, motels, hotels, nursing homes, schools, and hospitals. Use on wet-cleanable synthetic fibers. Do not use on wool. Vacuum carpet thoroughly prior to cleaning. Test fabric for colorfastness. Do not mix this product with other cleaning products. Follow the cleaning procedures specified by the manufacturer of the cleaning equipment. After using the product, set the carpet pile and protect the carpet from furniture legs and bases while drying. Do not over-wet. If applied to stain resistant nylon carpet, apply a fabric protector according to the manufacturer's directions

STORAGE AND DISPOSAL

(For Household/ Residential Use Products)

Store in original container in areas inaccessible to children. Nonrefillable container. Do not reuse or refill this container. Wrap container and put in trash or offer for recycling if available.

(For Commercial/Institutional/Industrial Uses)

Do not contaminate water, food or feed by storage or disposal.

Pesticide Storage: Open dumping is prohibited. Store in original container in areas inaccessible to children. Container Disposal: Nonrefillable container. Do not reuse or refill this container. Triple rinse (or equivalent). Offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

1166 012611

Sold By: Share Corporation • P.O. Box 245013 • Milwaukee, WI 53224 • 800-776-7192 • www. sharecorp.com

SHARE CORPORATION

CORPORATION		WORK HISTORY	
DESCRIPTION:	Viro-Stat RTU_QT	Vers.	Date
PRODUCT NUMBER:	1166	2	02.07.13
CUSTOMER:	Share Corporation		
SIZE:	5.0"W x 10.0"L		
APPLICATION:	Silkscreen		
COLOR(S):	PMS 234/280		
SIGNATURE:	BP		

ACCEPTED FOR REGISTRATION

Apr 12, 2013

New York State Department of Environmental Conservation Division of Materials Management Pesticide Product Registration

Doc id: 534935